arXiv:1611.01016v1 [math.QA] 3 Nov 2016

Differential and integral forms on non-commutative algebras

Tomasz Brzeziński

Mathematics Subject Classification (2010). Primary 58B30.
Keywords. Noncommutative geometry; differential forms; integral forms.

1. Introduction

These lectures describe an algebraic approach to differentiation and integration that is characteristic for non-commutative geometry. The material contained in Section 2 is standard and can be found in any text on non-commutative geometry, for example [4]. 2.5 and 2.6, which describe concepts introduced in [3], are exceptions. The bulk of Section 3 is based on [1] and [3], while the Berezin integral example is taken from [2].

2. Differential forms

2.1. **Differential graded algebras.** A *differential graded algebra* is a pair (Ω, d), where $\Omega = \oplus_{n\in\mathbb{Z}}\Omega^n$ is a graded algebra and $d : \Omega \to \Omega$ is a degree-one map that squares to zero and satisfies the graded Leibniz rule. Note that Ω^0 is an associative algebra and all the Ω^n are Ω^0-bimodules.

2.2. **Differential calculus.** Given an associative algebra A (over a field $\mathbb{K}$ of characteristic not 2), by a *differential calculus over* A we mean a differential graded algebra $(\Omega A, d)$, such that $\Omega^0 A = A$, ΩA is generated by $\Omega^1 A = Ad(A)$, and $\Omega^n A = 0$, for all $n < 0$. A calculus is said to be *N-dimensional*, if $\Omega^N A \neq 0$ and $\Omega^n A = 0$, for all $n > N$. The pair $(\Omega^1 A, d : A \to \Omega^1 A)$ is called a *first-order differential calculus*.

2.3. **Universal differential calculus.** Every algebra admits the *universal differential calculus*, defined as the tensor product algebra over the kernel of the multiplication map μ on A, with the exterior derivation $d : a \mapsto 1 \otimes a - a \otimes 1$, for all $a \in A$, and then extended to the whole of $T_A(\ker \mu)$ by the graded Leibniz rule.

Every first-order differential calculus $(\Omega^1 A, d)$ can by extended universally to the full calculus by defining ΩA as the quotient of the tensor algebra $T_A(\Omega^1 A)$ by the relations coming from the graded Leibniz rule and $d^2 = 0$.

2.4. **Volume form.** An N-dimensional calculus is said to *admit a volume form* if $\Omega^N A \cong A$ as a left and right A-module. Any free generator v of $\Omega^N A$ as a left and right A-module (if it exists) is called a *volume form*.

2.5. **Skew multi-derivations.** If $(\Omega A, d)$ is such that $\Omega^1 A$ is a finitely generated as a left A-module, then any left A-module basis $\{\omega_1, \dots, \omega_n\}$ of $\Omega^1 A$ induces maps $\partial_i, \sigma_{ij} : A \to A$, $i, j = 1, \dots, n$, by

$$da = \sum_i \partial_i(a)\omega_i, \qquad \omega_i a = \sum_j \sigma_{ij}(a)\omega_j. \tag{2.1}$$

These necessarily satisfy

$$\sigma_{ij}(1) = 1 \qquad \sigma_{ij}(ab) = \sum_k \sigma_{ik}(a)\sigma_{kj}(b), \tag{2.2a}$$

$$\partial_j(ab) = \sum_i \partial_i(a)\sigma_{ij}(b) + a\partial_j(b). \tag{2.2b}$$

A system $(\partial_j, \sigma_{ij})_{i,j=1}^n$ is called a *skew multi-derivation*. Any skew multi-derivation induces a calculus on A by formulae (2.1), provided there exist $a_i^{\alpha_i}, b_i^{\alpha_i} \in A$ such that $\sum_{\alpha_i} a_i^{\alpha_i}\partial_j(b_i^{\alpha_i}) = \delta_{ij}$. If such elements exist $(\partial_j, \sigma_{ij})_{i,j=1}^n$ is said to be *orthogonal*.

The conditions (2.2a) are equivalent to the statement that the map $\sigma : A \to M_n(A)$, $a \mapsto (\sigma_{ij}(a))_{i,j=1}^n$, where $M_n(A)$ denotes the ring of $n \times n$ matrices with entries from A, is an algebra homomorphism.

2.6. **Free multi-derivations.** A skew multi-derivation $(\partial_j, \sigma_{ij})_{i,j=1}^n$ is said to be *free*, provided there exist $\bar\sigma_{ij}, \hat\sigma_{ij} : A \to A$ that satisfy conditions (2.2a) and are such that, for all $a \in A$,

$$\sum_k \bar\sigma_{jk}(\sigma_{ik}(a)) = \sum_k \sigma_{kj}(\bar\sigma_{ki}(a)) = \sum_k \hat\sigma_{jk}(\bar\sigma_{ik}(a)) = \sum_k \bar\sigma_{kj}(\hat\sigma_{ki}(a)) = \delta_{ij}a.$$

If the matrix $\sigma = (\sigma_{ij})_{i,j=1}^n$ is triangular with invertible diagonal entries, then $(\partial_j, \sigma_{ij})_{i,j=1}^n$ is free.

2.7. **Diagonal and skew q-derivations.** If $\sigma = (\sigma_{ij})_{i,j=1}^n$ is diagonal, then relations (2.2b) separate into twisted Leibniz rules, $\partial_i(ab) = \partial_i(a)\sigma_{ii}(b) + a\partial_i(b)$, for all $i = 1, \dots, n$. Furthermore, if σ_{ii} is an invertible map and there exists $q_i \in \mathbb{K}$ such that $\sigma_{ii}^{-1} \circ \partial_i \circ \sigma_{ii} = q_i\partial_i$, then $(\partial_i, \sigma_{ii})$ is called a *skew q_i-derivation*.

2.8. **Calculus for q-polynomials.** A *q-polynomial algebra* or the *quantum plane* is the algebra $A = \mathbb{K}_q[x, y]$ generated by x, y subject to the relation $xy = qyx$. The elements of A are finite combinations of monomials $x^r y^s$.

The algebra A admits a first-order calculus freely generated by one-forms dx, dy and relations:

$$dxx = pxdx, \quad dyx = pq^{-1}xdy, \quad dxy = qydx + (p-1)xdy, \quad dyy = pydy,$$

where p is a non-zero scalar. It is understood that $d : x \mapsto dx, y \mapsto dy$. The universal extension of this calculus necessarily yields $dxdy = -qp^{-1}dydx$, $(dx)^2 = (dy)^2 = 0$, and it is a two-dimensional calculus with a volume form, e.g. $v = dxdy$.

Setting $\omega_1 = dx$, $\omega_2 = dy$ one easily finds that the associated skew multi-derivation $(\partial_i, \sigma_{ij})_{i,j=1}^2$ is free with the upper-triangular matrix-valued endomorphism

$$\sigma(x^r y^s) = \begin{pmatrix} p^r q^s x^r y^s & p^r(p^s - 1)x^{r+1}y^{s-1} \\ 0 & p^{r+s}q^{-r}x^r y^s \end{pmatrix}.$$

2.9. **Inner calculus.** A calculus $(\Omega A, d)$ is said to be *inner* if there exists $\theta \in \Omega^1 A$, such that $d(\omega) = \theta\omega - (-1)^n\omega$, for all $\omega \in \Omega^n A$. Note that $d^2(\omega) = 0$ implies that θ^2 is central in ΩA. Also θ satisfies the Cartan-Maurer equations $d\theta = 2\theta^2$.

As an example, consider a one-dimensional calculus on the Laurent polynomial ring $A = \mathbb{K}[x, x^{-1}]$, given by Jackson's q-derivation

$$\partial_q(f) = \frac{f(qx) - f(x)}{(q-1)x}, \tag{2.3}$$

where taking the limit is understood in case $q = 1$. If $q \neq 1$, this calculus is inner with $\theta = \frac{1}{q-1}x^{-1}dx$, otherwise it is not inner.

3. Integral forms

3.1. **Divergence.** Let $(\Omega A, d)$ be a differential calculus on an algebra A. We will denote by $\mathfrak{I}_n A$, the Abelian group of all right A-linear maps $\Omega^n A \to A$. For all $n \geq m$, consider maps

$$\cdot : \mathfrak{I}_n A \otimes \Omega^m A \to \mathfrak{I}_{n-m} A, \qquad f \otimes \omega \mapsto f \cdot \omega, \quad (f \cdot \omega)(\omega') = f(\omega\omega').$$

In particular, $\cdot$ makes $\mathfrak{I}_n A$ into a right A-module.

A *divergence* is a linear map $\nabla_0 : \mathfrak{I}_1 A \to A$, such that, for all $a \in A$, $\nabla_0(f \cdot a) = \nabla_0(f)a + f(da)$. A divergence is extended to a family of maps $\nabla_n : \mathfrak{I}_{n+1} A \to \mathfrak{I}_n A$, by $\nabla_n(f)(\omega) = \nabla_0(f \cdot \omega) + (-1)^{n+1} f(d\omega)$, for all $\omega \in \Omega^n A$.

The cokernel map $\Lambda : A \to A/\mathrm{Im}\nabla_0$ is called the *integral* associated to ∇_0.

3.2. **Integral forms.** A divergence ∇_0 is said to be *flat*, provided $\nabla_0 \circ \nabla_1 = 0$. It is then the case that, for all n, $\nabla_n \circ \nabla_{n+1} = 0$, and hence there is a complex, $\cdots \xrightarrow{\nabla_2} \mathfrak{I}_2 \xrightarrow{\nabla_1} \mathfrak{I}_1 \xrightarrow{\nabla_0} A$, known as the complex of *integral forms*.

3.3. **The inner case.** If $(\Omega A, d)$ is an inner differential calculus with the exterior derivative given by a graded commutator with $\theta \in \Omega^1 A$, then $\nabla_0 : \mathfrak{I}_1 A \to A$, $f \mapsto -f(\theta)$ is a divergence. One easily finds that $\nabla_1(f)(\omega) = f(\theta\omega)$, and so this divergence is flat, provided $\theta^2 = 0$.

3.4. **Divergences and multi-derivations.** Let $(\partial_i, \sigma_{ij}; \bar{\sigma}_{ij}, \hat{\sigma}_{ij})_{i,j=1}^n$ be a free skew multi-derivation on A, and let $(\Omega^1 A, d)$ be the associated first-order calculus with generators $\omega_1, \ldots, \omega_n$. Let $\xi_i \in \mathfrak{I}_i$ be the dual basis to the ω_i, i.e. the ξ_i are given by $\xi_i(\omega_j) = \delta_{ij}$. Then

$$\nabla_0 : \mathfrak{I}_1 A \to A, \qquad f \mapsto \sum_{i,j,k} \bar{\sigma}_{kj} \left(\partial_i \left(f \left(\hat{\sigma}_{ki} \left(\omega_i\right)\right)\right)\right), \tag{3.1}$$

is a unique divergence such that $\nabla_0(\xi_i) = 0$, for all $i = 1, \ldots, n$.

In particular, if σ is diagonal and all the $(\partial_i, \sigma_{ii})$ are skew q_i-derivations, then

$$\nabla_0(f) = \sum_i q_i \partial_i \left(f \left(\omega_i\right)\right).$$

3.5. **Cauchy's integral formula.** Let $A = \mathbb{K}[x, x^{-1}]$ be the Laurent polynomial ring with the one-dimensional calculus $(\Omega A, d)$ given by Jackson's q-derivative (2.3). In this case ∂_q is twisted by the automorphism $\sigma(f(x)) = f(qx)$, and (∂_q, σ) is a skew q-derivation. $\Omega^1 A$ is generated by dx, and hence the corresponding divergence (3.1) is $\nabla_0(f) = q\partial_q \left(f \left(dx\right)\right)$. For all $f \in \mathfrak{I}_1 A$, define $f_x \in \mathbb{K}[x, x^{-1}]$ by $f_x(x) := f(dx)$. Then

$$\nabla_0(f) = q \frac{f_x(qx) - f_x(x)}{(q-1)x}.$$

The image of ∇_0 consists of all of $\mathbb{K}[x, x^{-1}]$ except the monomials αx^{-1}. Therefore, the integral is

$$\Lambda : \mathbb{K}[x, x^{-1}] \to \mathbb{K}, \qquad a \mapsto \mathrm{res}(a)\Lambda(x^{-1}).$$

In case $\mathbb{K} = \mathbb{C}$ we can normalise Λ as $\Lambda(x^{-1}) = 2\pi i$, and obtain the Cauchy integral formula.

3.6. **Calculus on quantum groups.** If A is a coordinate algebra of a compact quantum group (over $\mathbb{C}$), then every left-covariant differential calculus gives rise to a free multi-derivation [5], and hence there is a canonical divergence ∇_0 (3.1) and the corresponding integral Λ. Any right integral λ on A (the Haar measure) factors uniquely through Λ, i.e. there exists unique $\varphi : A/\mathrm{Im}\nabla_0 \to \mathbb{C}$ such that $\lambda = \varphi \circ \Lambda$.

3.7. **Berezin's integral.** Let A be a superalgebra of (integrable) real functions on the supercircle $S^{1|1}$. That is, A consists of $a(x, \vartheta) = a^0(x) + a^1(x)\vartheta$, where $a^i : [0, 1] \to \mathbb{R}$ are (integrable) functions such that $a^i(0) = a^i(1)$ and ϑ is a Grassmann variable, $\vartheta^2 = 0$. The differentiation on A is defined by

$$\partial_x a(x, \vartheta) := \frac{da^0(x)}{dx} + \frac{da^1(x)}{dx}\vartheta, \qquad \partial_\vartheta a(x, \vartheta) := a^1(x).$$

One easily checks that $\partial = (\partial_x, \partial_\vartheta)$ is a free skew multi-derivation with the twisting matrix-valued endomorphism $\sigma(a(x, \vartheta)) = \begin{pmatrix} a(x, \vartheta) & 0 \\ 0 & a(x, -\vartheta) \end{pmatrix}$.

The calculus $\Omega^1 A$ is freely generated by dx and $d\vartheta$. The corresponding divergence 3.4 comes out as

$$\nabla_0(f)(x,\vartheta) = \partial_x f_x(x,\vartheta) - \partial_\vartheta f_\vartheta(x,\vartheta) = \frac{df^0_x(x)}{dx} - f^1_\vartheta(x) + \frac{df^0_\vartheta(x)}{dx}\vartheta,$$

where $f_x(x,\vartheta) := f(dx)(x,\vartheta) = f^0_x(x) + f^1_x(x)\vartheta$ and $f_\vartheta(x,\vartheta) := f(d\vartheta)(x,\vartheta) = f^0_\vartheta(x) + f^1_\vartheta(x)\vartheta$.

If $a(x,\vartheta)$ is purely even, i.e. $a(x,\vartheta) = a(x)$, then setting $f_x(x,\vartheta) = 0$ and $f_\vartheta(x,\vartheta) = -a(x)\vartheta$ we obtain $a(x) = \nabla_0(f)$. Thus, the integral Λ vanishes on the even part of A. Since Λ is the cokernel map of ∇_0, for all $f \in \mathfrak{I}_1 A$,

$$0 = \Lambda \circ \nabla_0(f) = \Lambda\left(\frac{df^0_x(x)}{dx} - f^1_\vartheta(x) + \frac{df^0_\vartheta(x)}{dx}\vartheta\right) = \Lambda\left(\frac{d}{dx} f^0_\vartheta(x)\vartheta\right).$$

On the other hand, $f^0_\vartheta(0) = f^0_\vartheta(1)$, so $\int_0^1 \frac{d}{dx} f^0_\vartheta(x)dx = 0$. By the universality of Λ, $\Lambda(a^1(x)\vartheta) = \int_0^1 a^1(x)dx$. Therefore,

$$\Lambda(a^0(x) + a^1(x)\vartheta) = \int_0^1 a^1(x)dx,$$

i.e. Λ is the Berezin integral on the supercircle.

Acknowledgment

I would like to thank the organizers and participants of the 2016 Białowieża School for creating very friendly and stimulating atmosphere.

References

[1] T. Brzeziński, *Non-commutative connections of the second kind.* J. Algebra Appl. **7** (2008), 557–573.

[2] T. Brzeziński, *Divergences on projective modules and non-commutative integrals.* Int. J. Geom. Methods Mod. Phys. **8** (2011), 885–896.

[3] T. Brzeziński, L. El Kaoutit and C. Lomp, *Non-commutative integral forms and twisted multi-derivations.* J. Noncommut. Geom. **4** (2010), 281–312.

[4] M. Dubois-Violette, *Lectures on graded differential algebras and noncommutative geometry.* Noncommutative differential geometry and its applications to physics (Shonan, 1999), pp. 245–306, Math. Phys. Stud. **23**, Kluwer, Dordrecht (2001).

[5] S.L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups).* Comm. Math. Phys. **122** (1989),125–170.

Tomasz Brzeziński
Department of Mathematics, Swansea University, Swansea SA2 8PP, U.K.
Department of Mathematics, University of Białystok, K. Ciołkowskiego 1M, 15-245 Białystok, Poland
e-mail: T.Brzezinski@swansea.ac.uk